

Tailwaters Studies Summary

Joint Fisheries Investigation Plan (JFIP)
NY & PA

Approach

ANGLERS

DELAWARE TAILWATERS

CREEL SURVEY PLAN

2018-2019

Prepared for the Fisheries Investigation Plan for the Delaware Tailwaters

February 28, 2018






Department of Environmental Conservation

Timothy Pokorny

NYSDEC Coldwater Fisheries Biologist Region 4 – Stamford, NY






Department of Environmental Conservation

Daryl Pierce

Pennsylvania Fish & Boat Commission Area 5 Biologist/Manager Bushkill, PA







JFIP Electrofishing









Young-of-the-year brown trout and rainbow trout collected from five West Branch sampling locations and nine sampling locations on tributaries to the West Branch, 2017 to 2020








Adult sized brown trout and rainbow trout from the nighttime boat electrofishing in the West Branch, 2017 to 2021.









Size distribution of brown trout and rainbow trout captured from the nighttime boat electrofishing in the West Branch, 2017 to 2021.








Estimated angler hours (N) within the Delaware tailwaters and the West Branch for 2018 and 2019 survey years compared to historical estimations from the 1990's Delaware tailwaters fishery.







Length class distribution (inches) of brown trout and rainbow trout for released (gray line), harvested (orange line) and total catch (blue line) for the estimated total catch from the Delaware tailwaters (i.e., East and West branches and Delaware River, combined).



Redd Counts: West and East Branches







Summary Statements

- Brown Trout populations abundant and stable in West Branch for adults and YOY
- Rainbow Trout populations low and stable in West Branch for adults and YOY
- The size distribution of both Brown and Rainbow Trout shows a high percent of fish in the 12-18" range
- Creel data shows that effort has doubled in comparison to fishing effort in the 1990s
- Comparisons between length distribution from creel and electrofishing data are very similar: validating utility of electrofishing data to assess fishery
- Spawning occurring in both East and West Branches at high rates

